

ON THE LAGRANGIAN ANGLE AND THE KÄHLER ANGLE OF IMMERSSED SURFACES IN THE COMPLEX PLANE $\mathbb{C}^2$

XINGXIAO LI* AND XIAO LI

ABSTRACT. In this paper, we discuss the Lagrangian angle and the Kähler angle of immersed surfaces in $\mathbb{C}^2$. Firstly, we provide an extension of Lagrangian angle, Maslov form and Maslov class to more general surfaces in $\mathbb{C}^2$ than Lagrangian surfaces, and then naturally extend a theorem by J.-M. Morvan to surfaces of constant Kähler angle, together with an application showing that the Maslov class of a compact self-shrinker surface with constant Kähler angle is generally non-vanishing. Secondly, we obtain two pinching results for the Kähler angle which imply rigidity theorems of self-shrinkers with Kähler angle under the condition that $\int_M |h|^2 e^{-\frac{|x|^2}{2}} dV_M < \infty$, where h and x denote, respectively, the second fundamental form and the position vector of the surface.

1. INTRODUCTION

Suppose that $(N, \mathbf{J})$ is an (almost) Hermitian manifold with ω its Kähler form. Then the Riemannian metric $\langle \cdot, \cdot \rangle$ on N is related with ω by

$$\langle U, V \rangle = \omega(U, \mathbf{J}V), \quad U, V \in TN. \quad (1.1)$$

Let $x : M^m \rightarrow N$ be an immersed submanifold with tangent space TM^m , normal space $T^\perp M^m$ and volume form dV_M of the induced metric. Then we call x to be totally real if $J(x_* TM^m) \subset T^\perp M^m$; if, furthermore, $\dim_{\mathbb{C}} N = m$, then x is called Lagrangian. In the special case of $m = 2$, we can introduce the concept of Kähler angle to define alternatively the totally real surfaces and Lagrangian surfaces. According to [9], the Kähler angle θ of an immersed surface $x : M^2 \rightarrow N$ is defined by

$$x^* \omega = \cos \theta dV_M, \quad \theta \in [0, \pi]. \quad (1.2)$$

Then x is totally real if and only if $\cos \theta \equiv 0$, or equivalently, the Kähler angle θ of x is identical to $\frac{\pi}{2}$; x is Lagrangian in N if and only if $\theta \equiv \frac{\pi}{2}$ and $\dim_{\mathbb{C}} N = 2$. By the way, if $\cos \theta > 0$ everywhere, then x is called symplectic.

The Kähler angle of conformal minimal surfaces in complex projective space $\mathbb{C}P^n$ has been extensively studied. For example, each of the maps in the Veronese sequence gives a minimal immersion of 2-sphere into $\mathbb{C}P^n$ that has a constant Kähler angle (see [3], [19], [21]). The Kähler angle of surfaces immersed in the nearly Kähler manifold $\mathbb{S}^6$ is studied, for example, in [17] and [16].

2000 *Mathematics Subject Classification.* Primary 53A30; Secondary 53B25.

Key words and phrases. Kähler angle, Lagrangian angle, self-shrinker.

Research supported by National Natural Science Foundation of China (No. 11171091, 11371018).

Now put $N = \mathbb{C}^m$, the complex Euclidean space with complex coordinates $(z^1, \dots, z^m)$. Let $\Omega = dz^1 \wedge \dots \wedge dz^m$ be the globally defined *holomorphic volume form*. For a Lagrangian submanifold $x : M^m \rightarrow \mathbb{C}^m$, the Lagrangian angle of x is by definition a multi-valued function $\beta_x : M^m \rightarrow \mathbb{R}/2\pi\mathbb{Z}$ given by

$$\Omega_M := x^*\Omega = e^{\sqrt{-1}\beta_x} dV_M.$$

Note that, although the Lagrangian angle β_x can not be determined globally in general, its gradient $\nabla\beta_x$ is clearly a well-defined vector field on M^m .

In [18], J.-M. Morvan proved an important formula by which the mean curvature and the Lagrangian angle of a Lagrangian submanifold are linked to each other. Precisely we have

Theorem 1.1 ([18]). *Let $x : M^m \rightarrow \mathbb{C}^m$ be a Lagrangian submanifold and $\mathbf{J}$ be the canonical complex structure of $\mathbb{C}^m$. Then the mean curvature vector H and the Lagrangian angle β_x meet the following formula:*

$$H = \mathbf{J}x_*(\nabla\beta_x), \quad (1.3)$$

where ∇ denotes the gradient operator on M^m .

Consequently, x is minimal if and only if its *Maslov form* $\alpha := -d\beta_x$ vanishes identically, implying that x has a trivial *Maslov class* $[\alpha] \in H^1(M)$, where $H^1(M)$ is the first homology group of M^m .

Remark 1.1. Formulas similar or related to (1.3) have also been obtained by some other authors, say, in [10], [20] and etc. In particular, A. Arsie has extended Theorem 1.1 in [2] to Lagrangian submanifolds in general Calabi-Yau manifolds.

Theorem 1.1 and its various extensions have many applications and citations in a lot of literatures, including papers on the Lagrangian mean curvature flow. As mentioned in [5], there is a direct interesting application of (1.3) first given by Smoczyk, and then by Castro and Lerman ([5]) using a different method: *No compact and orientable Lagrangian self-shrinkers to the Lagrangian mean curvature flow in $\mathbb{C}^m$ with trivial Maslov class*.

In this paper, we consider the Lagrangian angle and Kähler angle of immersed surfaces in $\mathbb{C}^2$. As the first part, we provide an extension of Lagrangian angle, Maslov form and Maslov class to more general surfaces in $\mathbb{C}^2$ than Lagrangian surfaces, and then naturally extend Theorem 1.1 of J.-M. Morvan to surfaces of constant Kähler angle, together with an application showing that the Maslov class of a compact self-shrinker surface with constant Kähler angle is generally non-vanishing. Precisely, we shall prove

Theorem 1.2. *Let $x : M^2 \rightarrow \mathbb{C}^2$ be an immersed surface with constant Kähler angle θ . Then the mean curvature vector H , the canonical complex structure $\mathbf{J}$ of $\mathbb{C}^2$ and the Lagrangian angle β_x meet the following formula:*

$$\sin^2 \theta x_*(\nabla\beta_x) = -(\mathbf{J}H)^\top, \quad (1.4)$$

where $^\top$ denotes the projection onto the tangent space $x_*(TM^2)$.

Clearly, if x is a Lagrangian surface then (1.4) agrees with (1.3).

Note that an immersion $x : M^m \rightarrow \mathbb{R}^{m+p}$ is called a *self-shrinker* (of the mean curvature flow) if $H = -x^\perp$, where $^\perp$ denotes the orthogonal projection to the normal bundle of x . Thus an application of Theorem 1.2 proves the following

Theorem 1.3. *Let $x : M^2 \rightarrow \mathbb{C}^2$ be a self-shrinker with constant Kähler angle $\theta \in (0, \pi)$. If M is compact and orientable, then the Maslov class $[\alpha] \neq 0$.*

Corollary 1.4. *There are no self-shrinkers with constant Kähler angle $\theta \in (0, \pi)$ in $\mathbb{C}^2$ with the topology of a sphere.*

As for symplectic surfaces there has been a fairly large amount of work. In particular, if the initial surface of a mean curvature flow in a Kähler-Einstein surface is symplectic, then the surface M_t at every time t is also symplectic ([6], [7]). So this kind of flow is referred to as a *symplectic mean curvature flow*. In [1], by using the self-adjoint property of a stability operator, Arezzo-Sun prove the following gap theorem (restated for the different definition of self-shrinkers here):

Theorem 1.5 ([1]). *Suppose that $x : M^2 \rightarrow \mathbb{C}^2$ is a complete symplectic self-shrinker with polynomial volume growth. If $|h|^2 \leq 2$, then $|h|^2 \equiv 0$ and $x(M^2)$ must be a plane.*

Moreover, Han-Sun prove a theorem in [13] that a translating soliton to the symplectic mean curvature flow in $\mathbb{C}^2$ with polynomial volume growth, non-positive normal curvature and bounded second fundamental form must be minimal in case that $\cos \theta \geq \delta > 0$. The following is a similar result for symplectic self-shrinkers:

Theorem 1.6 ([1]). *Let $x : M^2 \rightarrow \mathbb{C}^2$ be a complete symplectic self-shrinker with Kähler angle θ and polynomial volume growth. If $|h|^2$ is bounded and $\cos \theta \geq \delta > 0$, then $x(M^2)$ must be a plane.*

Recently Li-Wang proved the following theorem which can be viewed as a complement of the above theorems:

Theorem 1.7 ([14]). *Let $x : M^2 \rightarrow \mathbb{C}^2$ be a compact orientable Lagrangian self-shrinker, and h be the second fundamental form of x . If $|h|^2 \leq 2$, then $|h|^2 \equiv 2$ and $x(M^2)$ is the Clifford torus $S^1(1) \times S^1(1)$.*

Remark 1.2. Castro and Lerma also proved Theorem 1.7 in [5] under the additional condition that the Gauss curvature K of M^2 is either non-negative or non-positive.

To extend the above theorems, we provide in this paper the following main results:

Theorem 1.8. *Let $x : M^2 \rightarrow \mathbb{C}^2$ be a complete self-shrinker with Kähler angle θ satisfying $\cos \theta \geq 0$. If*

$$\int_M |h|^2 e^{-\frac{|x|^2}{2}} dV_M < \infty,$$

then θ is constant and $x(M^2)$ is either a Lagrangian surface or a plane.

To state the next main theorem, we need to introduce, as is done in [6] and [8], an almost complex structure $\mathbf{J}_M$ on $\mathbb{C}^2$ along x . Let $\{e_1, e_2\}$ be an oriented orthonormal frame field on M^2 of which $\{\omega^i\}$ is the dual frame, and $\{e_3, e_4\}$ be an oriented orthonormal normal frame field of x . Define

$$\mathbf{J}_M(x_*e_1) = x_*e_2, \quad \mathbf{J}_M(x_*e_2) = -x_*e_1, \quad \mathbf{J}_Me_3 = -e_4, \quad \mathbf{J}_Me_4 = e_3.$$

Then, if we write

$$h = \sum_{i,j} (h_{ij}^3 \omega^i \omega^j e_3 + h_{ij}^4 \omega^i \omega^j e_4),$$

then it holds that

$$\begin{aligned} (\bar{D}_{e_i} \mathbf{J}_M)(x_*e_1) &= \bar{D}_{e_i}(x_*e_2) - \mathbf{J}_M \bar{D}_{e_i}(x_*e_1) \\ &= -(h_{1i}^4 - h_{2i}^3)e_3 + (h_{2i}^4 + h_{1i}^3)e_4, \\ (\bar{D}_{e_i} \mathbf{J}_M)(x_*e_2) &= -\bar{D}_{e_i}(x_*e_1) - \mathbf{J}_M \bar{D}_{e_i}(x_*e_2) \\ &= -(h_{2i}^4 + h_{1i}^3)e_3 - (h_{1i}^4 - h_{2i}^3)e_4, \\ (\bar{D}_{e_i} \mathbf{J}_M)(e_3) &= -\bar{D}_{e_i}e_4 - \mathbf{J}_M \bar{D}_{e_i}e_3 \\ &= (h_{1i}^4 - h_{2i}^3)(x_*e_1) + (h_{2i}^4 + h_{1i}^3)(x_*e_2), \\ (\bar{D}_{e_i} \mathbf{J}_M)(e_4) &= \bar{D}_{e_i}e_3 - \mathbf{J}_M \bar{D}_{e_i}e_4, \\ &= -(h_{2i}^4 + h_{1i}^3)(x_*e_1) + (h_{1i}^4 - h_{2i}^3)(x_*e_2). \end{aligned}$$

Consequently, we have

$$|\bar{D}\mathbf{J}_M|^2 = 4 \sum_i ((h_{2i}^4 + h_{1i}^3)^2 + (h_{1i}^4 - h_{2i}^3)^2).$$

Note that $|\bar{D}\mathbf{J}_M|^2$ is globally defined on M^2 and thus is free of the choice of $\{e_1, e_2\}$ and $\{e_3, e_4\}$.

Theorem 1.9. *Let $x : M^2 \rightarrow \mathbb{C}^2$ be a complete self-shrinker with Kähler angle θ . If*

$$\int_M |h|^2 e^{-\frac{|x|^2}{2}} dV_M < \infty$$

and there exists a number $\lambda \in [0, 1)$ such that

$$|\nabla \theta|^2 \leq \frac{\lambda \cos^2 \theta |\bar{D}J_M|^2}{4(1 - \lambda \cos^2 \theta)}, \quad (1.5)$$

then θ is constant and $x(M^2)$ is either a Lagrangian surface or a plane.

Corollary 1.10. *Let $x : M^2 \rightarrow \mathbb{C}^2$ be a compact orientable self-shrinker with Kähler angle θ satisfying either $\cos \theta \geq 0$ or*

$$|\nabla \theta|^2 \leq \frac{\lambda \cos^2 \theta |\bar{D}J_M|^2}{4(1 - \lambda \cos^2 \theta)}$$

for some $\lambda \in [0, 1)$. If $|h|^2 \leq 2$, then

$$\theta = \frac{\pi}{2}, \quad |h|^2 = 2$$

and $x(M^2)$ is the Clifford torus $S^1(1) \times S^1(1)$.

Proof of Corollary 1.10: Since M^2 is compact, by Theorem 1.8 or Theorem 1.9, we know that $x : M^2 \rightarrow \mathbb{C}^2$ is a Lagrangian surface. Then it follows from Theorem 1.7 that $x(M^2)$ must be the Clifford torus $S^1(1) \times S^1(1)$. $\square$

2. KÄHLER ANGLES OF SURFACES AND SELF-SHRINKERS OF MEAN CURVATURE FLOW

In this section, we recall some necessary basics and formulas about the Kähler angle and self-shrinkers. First we make the following convention for the ranges of indices which are to be agreed with, if no other specification, throughout this present paper:

$$1 \leq i, j, k \leq 2, \quad 3 \leq a, b, c \leq 4, \quad 1 \leq A, B, C \leq 4.$$

On $\mathbb{C}^2$ we use the standard complex coordinates

$$z_1 = x_1 + \sqrt{-1}y_1, \quad z_2 = x_2 + \sqrt{-1}y_2.$$

The canonical complex structure $\mathbf{J}$ on $\mathbb{C}^2$ is given by

$$\mathbf{J} \frac{\partial}{\partial x_i} = \frac{\partial}{\partial y_i}, \quad \mathbf{J} \frac{\partial}{\partial y_i} = -\frac{\partial}{\partial x_i}.$$

Let $x : M^2 \rightarrow \mathbb{C}^2$ be an immersed surface and $\omega, \langle \cdot, \cdot \rangle$ be the Kähler form and the Riemannian metric on $\mathbb{C}^2$, respectively. The Kähler angle θ of x in $\mathbb{C}^2$ is defined by (1.2) which is equivalent to

$$\cos \theta = \langle \mathbf{J}(x_*e_1), x_*e_2 \rangle, \quad (2.1)$$

where $\{e_1, e_2\}$ is an oriented orthonormal frame field on M^2 .

Starting from any given $\{e_1, e_2\}$ on M^2 , we can construct along x an orthonormal frame field $\{x_*e_1, x_*e_2, e_3, e_4\}$ of $\mathbb{C}^2$, such that the following are satisfied in case $\theta \neq 0, \pi$ [17]:

$$\begin{aligned} \mathbf{J}(x_*e_1) &= (x_*e_2) \cos \theta + e_3 \sin \theta, & \mathbf{J}(x_*e_2) &= -(x_*e_1) \cos \theta + e_4 \sin \theta, \\ \mathbf{J}(e_3) &= -(x_*e_1) \sin \theta - e_4 \cos \theta, & \mathbf{J}(e_4) &= -(x_*e_2) \sin \theta + e_3 \cos \theta. \end{aligned} \quad (2.2)$$

Denote by $D, \bar{D}$ and $D^\perp$ the Levi-Civita connections on $M^2, \mathbb{C}^2$ and the normal bundle, respectively. Then the formulas of Gauss and Weingarten are given by

$$\bar{D}_X(x_*Y) = x_*(D_XY) + h(X, Y), \quad \bar{D}_X\xi = -x_*(A_\xi X) + D_X^\perp \xi,$$

where X and Y are tangent vector fields, ξ is a normal vector field on x , h denotes the second fundamental form and A is the shape operator.

In what follows we identify M^2 with $x(M^2)$ and omit x_* from some formulas and equations.

Let $\{\omega^A\}$ be the dual frame field of $\{e_A\}$, and $\{\omega_A^B\}$ the components of the Levi-Civita connection of $\mathbb{C}^2$ with respect to $\{e_A\}$. Then a direct computation similar to that in [17] using (2.2) shows

Lemma 2.1 (cf. [17]). *If $\sin \theta \neq 0$, then the following identities hold for an immersed surface $x : M^2 \rightarrow \mathbb{C}^2$:*

$$d\theta = \omega_1^4 - \omega_2^3, \quad (2.3)$$

$$(\omega_1^3 + \omega_2^4) \cos \theta + (\omega_3^4 - \omega_1^2) \sin \theta = 0. \quad (2.4)$$

Note that $\omega_i^a = \sum_j h_{ij}^a \omega^j$, where h_{ij}^a are components of the second fundamental form h of x . Then by (2.3), the Kähler angle θ of x is constant if and only if

$$h_{11}^4 = h_{12}^3, \quad h_{12}^4 = h_{22}^3. \quad (2.5)$$

Now we turn to self-shrinkers of the mean curvature flow in Euclidean space.

Let $x : M^m \rightarrow \mathbb{R}^{m+p}$ be an immersed n -dimensional submanifold in the $(m+p)$ -dimensional Euclidean space with the mean curvature vector H . Then x is called a *self-shrinker* (of the mean curvature flow) if $H = -x^\perp$, where $^\perp$ denotes the orthogonal projection to the normal bundle of x .

For a self-shrinker $x : M^m \rightarrow \mathbb{R}^{m+p}$, there is an important operator $\mathcal{L}$ acting on smooth functions that was first introduced and used by Colding and Minicozzi ([11]) to study self-shrinkers. The definition of $\mathcal{L}$ is as follows:

$$\mathcal{L} = \Delta - \langle x, \nabla \cdot \rangle = e^{\frac{|x|^2}{2}} \operatorname{div} (e^{-\frac{|x|^2}{2}} \nabla \cdot), \quad (2.6)$$

where Δ , ∇ and div denote the Laplacian, gradient and divergence on M^m , respectively.

Given an orthonormal tangent frame field $\{e_i; 1 \leq i \leq m\}$ on M^m with the dual $\{\omega^i\}$ and a normal frame field $\{e_\alpha; m+1 \leq \alpha \leq m+p\}$, write $h = \sum_{\alpha, i, j} h_{ij}^\alpha \omega^i \omega^j e_\alpha$. Then the mean curvature vector is by definition $H = \sum_\alpha H^\alpha e_\alpha$ with $H^\alpha = \sum_i h_{ii}^\alpha$.

The following lemma is fundamental for study of self-shrinkers.

Lemma 2.2 ([4]). *Let $x : M^m \rightarrow \mathbb{R}^{m+p}$ be a self-shrinker. Then*

$$H_{,i}^\alpha = \sum_j h_{ij}^\alpha \langle x, e_j \rangle. \quad (2.7)$$

We end this section with another lemma which is essential to our argument.

Lemma 2.3 ([15]). *Let $x : M^m \rightarrow \mathbb{R}^{m+p}$ be a complete immersed submanifold. If u and v are C^2 -smooth functions with*

$$\int_M (|u \nabla v| + |\nabla u \nabla v| + |u \mathcal{L} v|) e^{-\frac{|x|^2}{2}} dV_M < \infty,$$

then it holds that

$$\int_M u \mathcal{L} v e^{-\frac{|x|^2}{2}} dV_M = - \int_M \langle \nabla u, \nabla v \rangle e^{-\frac{|x|^2}{2}} dV_M.$$

3. THE LAGRANGIAN ANGLE FOR IMMERSED SURFACES IN $\mathbb{C}^2$

In this section, we first define a natural extension of the concept of Lagrangian angles to general immersed surfaces in $\mathbb{C}^2$, and then provide a proof of Theorem 1.2.

Given such a surface $x : M^2 \rightarrow \mathbb{C}^2$ with an orthonormal tangent frame field $\{e_1, e_2\}$ and its dual $\{\omega^1, \omega^2\}$. As mentioned in the introduction, let Ω denote the holomorphic volume form on $\mathbb{C}^2$ given by $\Omega = dz_1 \wedge dz_2$, where $z_1 = x_1 + \sqrt{-1}y_1$, $z_2 = x_2 + \sqrt{-1}y_2$.

Lemma 3.1. *When pulling back by x , Ω is a complex multiple of the volume form dV_M . More precisely, there exists a complex function η_x such that $x^*\Omega = \eta_x dV_M$.*

Proof. In fact, $\eta_x = x^*\Omega(e_1, e_2) = \Omega(x_*e_1, x_*e_2)$ since $dV_M = \omega^1 \wedge \omega^2$. $\square$

Therefore, when $x^*\Omega \neq 0$, or the same, $\eta_x \neq 0$, there determined (up to a multiple of 2π) is a multi-valued angle function β_x , usually called the argument of η_x , by $\eta_x = |\eta_x|e^{\sqrt{-1}\beta_x}$.

Thus the following definition is natural:

Definition 3.1. Let $x : M^2 \rightarrow \mathbb{C}^2$ be an immersed surface with $x^*\Omega \neq 0$. Then the multi-valued angle function β_x is called the Lagrangian angle of x .

Remark 3.1. Clearly, $\alpha := -d\beta_x$ is a well defined closed 1-form on M^2 which is called the Maslov form of x and represents a cohomology class $[\alpha] \in H^1(M^2)$, referred to as the Maslov class of x .

Next we are to find how the Kähler angle θ influences the Lagrangian angle β_x for an immersion $x : M^2 \rightarrow \mathbb{C}^2$. To this end, we need the following simple lemma:

Lemma 3.2. *For any $A \in SU(2)$, it holds that $\eta_{Ax} = \eta_x$. In particular, $\beta_{Ax} = \beta_x$ in case that $\eta_x \neq 0$.*

Proof. Note that

$$SU(2) = \left\{ \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \mid a, b \in \mathbb{C}, |a|^2 + |b|^2 = 1 \right\}.$$

Write $z = (z_1, z_2) \in \mathbb{C}^2$ and, for any $A = \begin{pmatrix} a & b \\ -\bar{b} & \bar{a} \end{pmatrix} \in SU(2)$, $\tilde{z} \equiv Az = (\tilde{z}_1, \tilde{z}_2)$. Since $\det A = 1$, we find

$$A^*\Omega = d\tilde{z}_1 \wedge d\tilde{z}_2 = (\det A)dz_1 \wedge dz_2 = \Omega. \quad (3.1)$$

On the other hand, by Lemma 3.1, we have

$$x^*\Omega = \eta_x dV_M, \quad (Ax)^*\Omega = \eta_{Ax} dV_M.$$

Hence by (3.1)

$$\eta_{Ax} = ((Ax)^*\Omega)(e_1, e_2) = (x^*(A^*\Omega))(e_1, e_2) = (x^*\Omega)(e_1, e_2) = \eta_x.$$

$\square$

Now we suppose that the Kähler angle $\theta \in (0, \pi)$. For any $p \in M^2$, we can find an $A \in SU(2)$ to rotate $\mathbb{C}^2$ such that $(Ax)_*e_1 = \frac{\partial}{\partial x_1}$ at p . Then, since $(Ax)_*e_1$ and $(Ax)_*e_2$ are orthogonal, we obtain from (2.1) that

$$(Ax)_*e_2 = \cos \theta \frac{\partial}{\partial y_1} + \sin \theta (\cos \beta_{Ax} \frac{\partial}{\partial x_2} + \sin \beta_{Ax} \frac{\partial}{\partial y_2}).$$

Since

$$\Omega = (dx_1 + \sqrt{-1}dy_1) \wedge (dx_2 + \sqrt{-1}dy_2), \quad (3.2)$$

we compute

$$\Omega((Ax)_*e_1, (Ax)_*e_2) = \sin \theta (\cos \beta_{Ax} + \sqrt{-1} \sin \beta_{Ax}) = \sin \theta e^{\sqrt{-1}\beta_{Ax}}.$$

Hence

$$(Ax)^*\Omega = \sin \theta e^{\sqrt{-1}\beta_{Ax}} dV_M, \quad \text{at } p. \quad (3.3)$$

According to Lemma 3.2, we can get

$$x^*\Omega = \sin \theta e^{\sqrt{-1}\beta_x} dV_M, \quad \text{at } p. \quad (3.4)$$

Thus we have proved

Proposition 3.3. *Let $x : M^2 \rightarrow \mathbb{C}^2$ be an immersed surface with Kähler angle $\theta \in (0, \pi)$. Then $|\eta_x|^2 = \sin^2 \theta$. In particular, $\eta_x \neq 0$ everywhere on M^2 .*

We are in a position to give a proof of Theorem 1.2.

First we assume that $\sin \theta \neq 0$.

Note that, for any orthonormal frame field $\{e_1, e_2\}$, we can find a unique normal frame $\{e_3, e_4\}$ by (2.2). So, along $x(M^2)$, we have an orthonormal frame $\{x_*e_i, e_a\}$ of $\mathbb{C}^2$ with the dual frame field $\{\omega^A, 1 \leq A \leq 4\}$. As the first step to prove Theorem 1.2, we claim that

$$\Omega \circ x = \frac{1}{\sin \theta} e^{\sqrt{-1}\beta_x} (\omega^1 - \sqrt{-1}\mathbf{J}\omega^1) \wedge (\omega^2 - \sqrt{-1}\mathbf{J}\omega^2). \quad (3.5)$$

In fact, since Ω is a $(2, 0)$ -form and $\omega^1 - \sqrt{-1}\mathbf{J}\omega^1, \omega^2 - \sqrt{-1}\mathbf{J}\omega^2$ are two linearly independent $(1, 0)$ -forms, we can write

$$\Omega \circ x = \mu (\omega^1 - \sqrt{-1}\mathbf{J}\omega^1) \wedge (\omega^2 - \sqrt{-1}\mathbf{J}\omega^2)$$

for some complex-valued function μ on M^2 .

Since by (2.2)

$$\begin{aligned} \mathbf{J}(\omega^1) &= -\omega^2 \cos \theta - \omega^3 \sin \theta, & \mathbf{J}(\omega^2) &= \omega^1 \cos \theta - \omega^4 \sin \theta, \\ \mathbf{J}(\omega^3) &= \omega^1 \sin \theta + \omega^4 \cos \theta, & \mathbf{J}(\omega^4) &= \omega^2 \sin \theta - \omega^3 \cos \theta, \end{aligned} \quad (3.6)$$

it follows that

$$\frac{1}{2}(\omega^i - \sqrt{-1}\mathbf{J}\omega^i)(x_*e_i - \sqrt{-1}\mathbf{J}(x_*e_i)) = 1, \quad i = 1, 2; \quad (3.7)$$

$$\begin{aligned} & \frac{1}{2}(\omega^1 - \sqrt{-1}\mathbf{J}\omega^1)(x_*e_2 - \sqrt{-1}\mathbf{J}(x_*e_2)) \\ &= -\frac{1}{2}(\omega^2 - \sqrt{-1}\mathbf{J}\omega^2)(x_*e_1 - \sqrt{-1}\mathbf{J}(x_*e_1)) \\ &= \sqrt{-1} \cos \theta. \end{aligned} \quad (3.8)$$

So that

$$\Omega \left(\frac{x_*e_1 - \sqrt{-1}\mathbf{J}(x_*e_1)}{2}, \frac{x_*e_2 - \sqrt{-1}\mathbf{J}(x_*e_2)}{2} \right) = \mu \sin^2 \theta. \quad (3.9)$$

On the other hand, for any given $p \in M^2$, we can assume that $x_{*p}e_1 = \frac{\partial}{\partial x_1}|_{x(p)}$ because $A^*\Omega = \Omega$ for any $A \in SU(2)$. Then it holds that

$$x_*e_1 - \sqrt{-1}\mathbf{J}(x_*e_1) = \frac{\partial}{\partial x_1} - \sqrt{-1}\frac{\partial}{\partial y_1}, \quad (3.10)$$

$$\begin{aligned} x_*e_2 - \sqrt{-1}\mathbf{J}(x_*e_2) &= \cos \theta \frac{\partial}{\partial y_1} + \sin \theta \left(\cos \beta_x \frac{\partial}{\partial x_2} + \sin \beta_x \frac{\partial}{\partial y_2} \right) \\ &\quad + \sqrt{-1} \cos \theta \frac{\partial}{\partial x_1} - \sqrt{-1} \sin \theta \left(\cos \beta_x \frac{\partial}{\partial y_2} - \sin \beta_x \frac{\partial}{\partial x_2} \right) \end{aligned} \quad (3.11)$$

at p . Substitute (3.10) and (3.11) into (3.2) and obtain

$$\Omega \left(\frac{x_*e_1 - \sqrt{-1}\mathbf{J}(x_*e_1)}{2}, \frac{x_*e_2 - \sqrt{-1}\mathbf{J}(x_*e_2)}{2} \right) = \sin \theta e^{\sqrt{-1}\beta_x} \quad \text{at } p. \quad (3.12)$$

Comparing (3.9) with (3.12), we obtain

$$\mu(p) = \frac{1}{\sin \theta} e^{\sqrt{-1}\beta_x(p)}.$$

Thus (3.5) is proved by the arbitrariness of the point p .

The second step is to make use of the fact that Ω is parallel on $\mathbb{C}^2$. For any tangent vector field $X \in TM^2$, we have

$$\bar{D}_X \Omega = 0. \quad (3.13)$$

Note that the Kähler angle θ is constant. (3.13) is equivalent to

$$\begin{aligned} 0 &= \sqrt{-1} \sin^2 \theta d\beta_x(X) \Omega + \sin \theta e^{\sqrt{-1}\beta_x} (\bar{D}_X \omega^1 - \sqrt{-1} \bar{D}_X \mathbf{J} \omega^1) \wedge (\omega^2 - \sqrt{-1} \mathbf{J} \omega^2) \\ &\quad + \sin \theta e^{\sqrt{-1}\beta_x} (\omega^1 - \sqrt{-1} \mathbf{J} \omega^1) \wedge (\bar{D}_X \omega^2 - \sqrt{-1} \bar{D}_X \mathbf{J} \omega^2). \end{aligned} \quad (3.14)$$

Since the Levi-Civita connection $\bar{D}$ is also a complex one, that is, $\bar{D}\mathbf{J} = \mathbf{J}\bar{D}$, we know that $\bar{D}_X \omega^1 - \sqrt{-1} \bar{D}_X \mathbf{J} \omega^1$ is a linear combination of

$$\omega^1 - \sqrt{-1} \mathbf{J} \omega^1 \quad \text{and} \quad \omega^2 - \sqrt{-1} \mathbf{J} \omega^2,$$

or

$$\bar{D}_X \omega^1 - \sqrt{-1} \bar{D}_X \mathbf{J} \omega^1 = \rho_1 (\omega^1 - \sqrt{-1} \mathbf{J} \omega^1) + \rho_2 (\omega^2 - \sqrt{-1} \mathbf{J} \omega^2) \quad (3.15)$$

for two complex-valued functions ρ_1 and ρ_2 .

Moreover, from $\bar{D}\mathbf{J} = \mathbf{J}\bar{D}$ and (3.6), we have

$$\begin{aligned}\omega^1(\bar{D}_X\mathbf{J}e_1) &= \omega^1(\mathbf{J}(\bar{D}_Xe_1)) = (\mathbf{J}\omega^1)(\bar{D}_Xe_1) \\ &= (-\omega^2 \cos \theta - \omega^3 \sin \theta)(\bar{D}_Xe_1) \\ &= -\cos \theta \langle \bar{D}_Xe_1, e_2 \rangle - \sin \theta \langle \bar{D}_Xe_1, e_3 \rangle, \\ \omega^1(\bar{D}_X\mathbf{J}e_2) &= \omega^1(\mathbf{J}(\bar{D}_Xe_2)) = (\mathbf{J}\omega^1)(\bar{D}_Xe_2) \\ &= (-\omega^2 \cos \theta - \omega^3 \sin \theta)(\bar{D}_Xe_2) = -\sin \theta \langle \bar{D}_Xe_2, e_3 \rangle.\end{aligned}$$

From (3.7), (3.8) and (3.15) it follows that

$$\begin{aligned}\rho_1 - \sqrt{-1}\rho_2 \cos \theta &= (\bar{D}_X\omega^1 - \sqrt{-1}\bar{D}_X\mathbf{J}\omega^1)((e_1 - \sqrt{-1}\mathbf{J}e_1)/2) \\ &= -\sqrt{-1}(\cos \theta \langle \bar{D}_Xe_1, e_2 \rangle + \sin \theta \langle \bar{D}_Xe_1, e_3 \rangle).\end{aligned}\tag{3.16}$$

$$\begin{aligned}\sqrt{-1}\rho_1 \cos \theta + \rho_2 &= (\bar{D}_X\omega^1 - \sqrt{-1}\bar{D}_X\mathbf{J}\omega^1)((e_2 - \sqrt{-1}\mathbf{J}e_2)/2) \\ &= -\sqrt{-1} \sin \theta \langle \bar{D}_Xe_2, e_3 \rangle - \langle \bar{D}_Xe_2, e_1 \rangle.\end{aligned}\tag{3.17}$$

Combining (3.16) and (3.17) we can find ρ_1 and ρ_2 to obtain

$$\begin{aligned}&\sin \theta (\bar{D}_X\omega^1 - \sqrt{-1}\bar{D}_X\mathbf{J}\omega^1) \\ &= (\cos \theta \langle \bar{D}_Xe_2, e_3 \rangle - \sqrt{-1} \langle \bar{D}_Xe_1, e_3 \rangle) (\omega^1 - \sqrt{-1}\mathbf{J}\omega^1) \\ &\quad - (\sin \theta \langle \bar{D}_Xe_2, e_1 \rangle + \sqrt{-1} \langle \bar{D}_Xe_2, e_3 \rangle + \cos \theta \langle \bar{D}_Xe_1, e_3 \rangle) (\omega^2 - \sqrt{-1}\mathbf{J}\omega^2)\end{aligned}\tag{3.18}$$

Similarly we have

$$\begin{aligned}&\sin \theta (\bar{D}_X\omega^2 - \sqrt{-1}\bar{D}_X\mathbf{J}\omega^2) \\ &= (-\cos \theta \langle \bar{D}_Xe_1, e_4 \rangle - \sqrt{-1} \langle \bar{D}_Xe_2, e_4 \rangle) (\omega^2 - \sqrt{-1}\mathbf{J}\omega^2) \\ &\quad - (\sin \theta \langle \bar{D}_Xe_1, e_2 \rangle - \cos \theta \langle \bar{D}_Xe_2, e_4 \rangle + \sqrt{-1} \langle \bar{D}_Xe_1, e_4 \rangle) (\omega^1 - \sqrt{-1}\mathbf{J}\omega^1).\end{aligned}\tag{3.19}$$

Therefore we can rewrite (3.14) as

$$\begin{aligned}0 &= (\sqrt{-1} \sin \theta d\beta_x(X) + \cos \theta \langle \bar{D}_Xe_2, e_3 \rangle - \sqrt{-1} \langle \bar{D}_Xe_1, e_3 \rangle \\ &\quad - \cos \theta \langle \bar{D}_Xe_1, e_4 \rangle - \sqrt{-1} \langle \bar{D}_Xe_2, e_4 \rangle) \Omega.\end{aligned}\tag{3.20}$$

Using Gauss formula and (3.20), we find

$$\begin{aligned}\sqrt{-1} \sin \theta d\beta_x(X) &= -\cos \theta \langle \bar{D}_Xe_2, e_3 \rangle + \sqrt{-1} \langle \bar{D}_Xe_1, e_3 \rangle \\ &\quad + \cos \theta \langle \bar{D}_Xe_1, e_4 \rangle + \sqrt{-1} \langle \bar{D}_Xe_2, e_4 \rangle \\ &= -\cos \theta \langle h(X, e_2), e_3 \rangle + \sqrt{-1} \langle h(X, e_1), e_3 \rangle \\ &\quad + \cos \theta \langle h(X, e_1), e_4 \rangle + \sqrt{-1} \langle h(X, e_2), e_4 \rangle \\ &= -\cos \theta \langle \bar{D}_{e_2}X, e_3 \rangle + \sqrt{-1} \langle \bar{D}_{e_1}X, e_3 \rangle \\ &\quad + \cos \theta \langle \bar{D}_{e_1}X, e_4 \rangle + \sqrt{-1} \langle \bar{D}_{e_2}X, e_4 \rangle \\ &= \cos \theta \langle \bar{D}_{e_2}e_3, X \rangle - \sqrt{-1} \langle \bar{D}_{e_1}e_3, X \rangle \\ &\quad - \cos \theta \langle \bar{D}_{e_1}e_4, X \rangle - \sqrt{-1} \langle \bar{D}_{e_2}e_4, X \rangle\end{aligned}\tag{3.21}$$

But from (2.2) it is easily seen that

$$e_3 = \frac{1}{\sin \theta} \mathbf{J}e_1 - \frac{\cos \theta}{\sin \theta} e_2, \quad e_4 = \frac{1}{\sin \theta} \mathbf{J}e_2 + \frac{\cos \theta}{\sin \theta} e_1. \quad (3.22)$$

Inserting (3.22) to (3.21), we obtain

$$\begin{aligned} \sqrt{-1} \sin^2 \theta d\beta_x(X) &= \cos \theta \langle \bar{D}_{e_2}(\mathbf{J}e_1), X \rangle - \cos \theta \langle \bar{D}_{e_2}(\cos \theta e_2), X \rangle \\ &\quad - \sqrt{-1} \langle \bar{D}_{e_1}(\mathbf{J}e_1), X \rangle + \sqrt{-1} \langle \bar{D}_{e_1}(\cos \theta e_2), X \rangle \\ &\quad - \cos \theta \langle \bar{D}_{e_1}(\mathbf{J}e_2), X \rangle - \cos \theta \langle \bar{D}_{e_1}(\cos \theta e_1), X \rangle \\ &\quad - \sqrt{-1} \langle \bar{D}_{e_2}(\mathbf{J}e_2), X \rangle - \sqrt{-1} \langle \bar{D}_{e_2}(\cos \theta e_1), X \rangle \\ &= \cos \theta \langle \mathbf{J}D_{e_2}e_1, X \rangle + \cos \theta \langle \mathbf{J}h(e_2, e_1), X \rangle - \cos^2 \theta \langle D_{e_2}e_2, X \rangle \\ &\quad - \sqrt{-1} \langle \mathbf{J}D_{e_1}e_1, X \rangle - \sqrt{-1} \langle \mathbf{J}h(e_1, e_1), X \rangle + \sqrt{-1} \cos \theta \langle D_{e_1}e_2, X \rangle \\ &\quad - \cos \theta \langle \mathbf{J}D_{e_1}e_2, X \rangle - \cos \theta \langle \mathbf{J}h(e_1, e_2), X \rangle - \cos^2 \theta \langle D_{e_1}e_1, X \rangle \\ &\quad - \sqrt{-1} \langle \mathbf{J}D_{e_2}e_2, X \rangle - \sqrt{-1} \langle \mathbf{J}h(e_2, e_2), X \rangle - \sqrt{-1} \cos \theta \langle D_{e_2}e_1, X \rangle. \end{aligned}$$

For any given point $p \in M^2$, choose proper frame field $\{e_i\}$ satisfying $D_{e_i}e_j = 0$ at p . Then

$$\begin{aligned} \sin^2 \theta d\beta_x(X) &= - \langle \mathbf{J}(h(e_1, e_1)), X \rangle - \langle \mathbf{J}(h(e_2, e_2)), X \rangle \\ &= - \langle \mathbf{J}(h(e_1, e_1) + h(e_2, e_2)), X \rangle \\ &= - \langle (\mathbf{J}H)^\top, X \rangle \quad \text{at } p. \end{aligned}$$

Since both $d\beta_x(X)$ and $\langle (\mathbf{J}H)^\top, X \rangle$ are globally defined functions and thus free of the choice of the frame field $\{e_i\}$, and X is an arbitrary vector field on M^2 , we have $\sin^2 \theta \nabla \beta_x = -(\mathbf{J}H)^\top$ and Theorem 1.2 is proved for $\sin \theta \neq 0$.

When $\sin \theta = 0$, both sides of (1.4) vanish identically since any holomorphic curve in $\mathbb{C}^2$ is necessarily minimal. $\square$

4. EXAMPLES

In this section, we provide three examples of surfaces in $\mathbb{C}^2$. Example 4.1 and 4.2 are quoted from lecture notes by Jason Lotay.

Example 4.1. Let $x : \mathbb{R}^2 \rightarrow \mathbb{C}^2$ be given by

$$x(u_1, u_2) = (e^{\sqrt{-1}u_1}, e^{\sqrt{-1}u_2}), \quad (u_1, u_2) \in \mathbb{R}^2.$$

A short calculation shows that the immersion is an immersed Lagrangian self-shrinker with the image $x(\mathbb{R}^2) = S^1(1) \times S^1(1)$, a standard torus. Furthermore, if we denote $e_1 = \frac{\partial}{\partial u_1}$, $e_2 = \frac{\partial}{\partial u_2}$, then

$$x_*e_1 = (\sqrt{-1}e^{\sqrt{-1}u_1}, 0), \quad x_*e_2 = (0, \sqrt{-1}e^{\sqrt{-1}u_2}).$$

It is easy to check that

$$(x^*\Omega)(e_1, e_2) = e^{\sqrt{-1}(u_1+u_2+\pi)}$$

implying that the Lagrangian angle of x is given by $\beta_x = u_1 + u_2 + \pi$ up to multiples of 2π . Obviously, the Maslov class $[-d\beta_x]$ of x is nontrivial and, by Theorem 1.2 x is not minimal.

Example 4.2. Let $x : \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{C}^2$ be given by $(z_1, z_2) = x(u_1, u_2)$ with

$$z_1 = u_1 + \frac{u_1}{u_1^2 + u_2^2} \sqrt{-1}, \quad z_2 = u_2 + \frac{u_2}{u_1^2 + u_2^2} \sqrt{-1}, \quad (u_1, u_2) \in \mathbb{R}^2.$$

This is known as the Lagrangian catenoid and, with respect to the induced metric g , an orthonormal frame field can be chosen as

$$e_1 = \frac{u_1^2 + u_2^2}{\sqrt{1 + (u_1^2 + u_2^2)^2}} \frac{\partial}{\partial u_1}, \quad e_2 = \frac{u_1^2 + u_2^2}{\sqrt{1 + (u_1^2 + u_2^2)^2}} \frac{\partial}{\partial u_2}.$$

Then

$$dV_M = \frac{1 + (u_1^2 + u_2^2)^2}{(u_1^2 + u_2^2)^2} du_1 \wedge du_2, \quad \text{where } M^2 = (\mathbb{R}^2, g),$$

and

$$\begin{aligned} x_* e_1 &= \frac{u_1^2 + u_2^2}{\sqrt{1 + (u_1^2 + u_2^2)^2}} \left(1, 0, \frac{u_2^2 - u_1^2}{(u_1^2 + u_2^2)^2}, \frac{-2u_1 u_2}{(u_1^2 + u_2^2)^2} \right), \\ x_* e_2 &= \frac{u_1^2 + u_2^2}{\sqrt{1 + (u_1^2 + u_2^2)^2}} \left(0, 1, \frac{-2u_1 u_2}{(u_1^2 + u_2^2)^2}, \frac{u_1^2 - u_2^2}{(u_1^2 + u_2^2)^2} \right). \end{aligned}$$

A direct computation shows that

$$x^* \Omega = x^*(dz_1 \wedge dz_2) = \frac{1 + (u_1^2 + u_2^2)^2}{(u_1^2 + u_2^2)^2} du_1 \wedge du_2 = dV_M.$$

Thus

$$\eta_x \equiv 1 \quad \text{and} \quad \text{Arg}(\eta_x) = 2k\pi, \quad k \in \mathbb{Z}.$$

Therefore, the Lagrangian angle $\beta_x = 0$ up to multiples of 2π . Consequently by Theorem 1.2, x is a minimal immersion.

Example 4.3. For any given two real constants θ_1 and θ_2 , denote $\theta := \theta_1 + \theta_2$. Let $x : \mathbb{R}^2 \rightarrow \mathbb{C}^2$ be defined by $(z_1, z_2) = x(u_1, u_2)$ with

$$z_1 = u_1 \cos \theta_1 + \sqrt{-1} u_2 \cos \theta_2, \quad z_2 = -u_2 \sin \theta_2 - \sqrt{-1} u_1 \sin \theta_1, \quad (u_1, u_2) \in \mathbb{R}^2.$$

Choose $e_1 = \frac{\partial}{\partial u_1}$ and $e_2 = \frac{\partial}{\partial u_2}$. Then

$$x_* e_1 = (\cos \theta_1, 0, 0, -\sin \theta_1), \quad x_* e_2 = (0, -\sin \theta_2, \cos \theta_2, 0). \quad (4.1)$$

Thus

$$\mathbf{J}(x_* e_1) = (0, \sin \theta_1, \cos \theta_1, 0), \quad \mathbf{J}(x_* e_2) = (-\cos \theta_2, 0, 0, -\sin \theta_2). \quad (4.2)$$

It follows that

$$\langle \mathbf{J}(x_* e_1), x_* e_2 \rangle = \cos(\theta_1 + \theta_2) = \cos \theta,$$

implying that x is of constant Kähler angle θ .

On the other hand, since

$$\begin{aligned} x^*\Omega &= x^*(dz_1 \wedge dz_2) = (\cos \theta_1 du_1 + \sqrt{-1} \cos \theta_2 du_2) \wedge (-\sin \theta_2 du_2 - \sqrt{-1} \sin \theta_1 du_1) \\ &= -\sin(\theta_1 + \theta_2) du_1 \wedge du_2 = -\sin(\theta_1 + \theta_2) dV_{\mathbb{R}}, \end{aligned}$$

we know that the Lagrangian angle of x is $\beta_x = \pi$ up to multiples of 2π . So by Theorem 1.2, $H = 0$. Clearly, $x(\mathbb{R}^2)$ is actually a 2-plane in $\mathbb{C}^2$ and is indeed totally geodesic.

5. PROOF OF THE OTHER MAIN THEOREMS

In this section, we give proofs of other main theorems.

(1) *Proof of Theorem 1.3.*

For any tangent vector $v \in TM^2$, by the definition of a self-shrinker and the fundamental equations we find

$$-A_H v + D_v^\perp H = \bar{D}_v H = \bar{D}_v(-x^\perp) = \bar{D}_v(x^\top - x) = D_v x^\top + h(v, x^\top) - v.$$

So that

$$A_H v = v - D_v x^\top, \quad D_v^\perp H = h(v, x^\top). \quad (5.1)$$

For a given orthonormal frame field $\{e_1, e_2\}$ on M^2 , let the normal frame $\{e_3, e_4\}$ be defined via (2.2). Then a direct computation shows that

$$\mathbf{J}H = -\sin \theta (H^3 x_* e_1 + H^4 x_* e_2) + \cos \theta (H^4 e_3 - H^3 e_4) \quad (5.2)$$

which implies that

$$(\mathbf{J}H)^\top = -\sin \theta (H^3 x_* e_1 + H^4 x_* e_2), \quad (5.3)$$

$$(\mathbf{J}H)^\perp = \cos \theta (H^4 e_3 - H^3 e_4). \quad (5.4)$$

Since the Kähler angle θ is constant, by Lemma 2.1 and (2.5), we know that

$$h_{11}^4 = h_{12}^3, \quad h_{12}^4 = h_{22}^3, \quad \omega_1^2 = \omega_3^4 + \cot \theta (\omega_1^3 + \omega_2^4). \quad (5.5)$$

Applying (5.3) – (5.5), we have

$$\begin{aligned} D_{e_i}(\mathbf{J}H)^\top &= -\sin \theta D_{e_i}(H^3 e_1 + H^4 e_2) \\ &= -\sin \theta (e_i(H^3) e_1 + e_i(H^4) e_2 + H^3 \omega_1^2(e_i) e_2 + H^4 \omega_2^1(e_i) e_1) \\ &= -\sin \theta (e_i(H^3) e_1 + e_i(H^4) e_2 + (H^3 e_2 - H^4 e_1) \omega_1^2(e_i)) \\ &= -\sin \theta (e_i(H^3) e_1 + e_i(H^4) e_2 + (H^3 e_2 - H^4 e_1) \omega_3^4(e_i) \\ &\quad + \cot \theta (H^3 e_2 - H^4 e_1) (\omega_1^3 + \omega_2^4)(e_i)) \\ &= -\sin \theta (e_i(H^3) e_1 + e_i(H^4) e_2 + (H^3 e_2 - H^4 e_1) \omega_3^4(e_i)) \\ &\quad - \cos \theta (H^3 e_2 - H^4 e_1) (h_{1i}^3 + h_{2i}^4); \end{aligned}$$

and

$$\begin{aligned}
\mathbf{J}D_{e_i}^\perp H &= \mathbf{J}D_{e_i}^\perp (H^3 e_3 + H^4 e_4) \\
&= \mathbf{J}(e_i(H^3)e_3 + e_i(H^4)e_4 + H^3 D_{e_i}^\perp e_3 + H^4 D_{e_i}^\perp e_4) \\
&= \mathbf{J}(e_i(H^3)e_3 + e_i(H^4)e_4 + H^3 \omega_3^4(e_i)e_4 + H^4 \omega_4^3(e_i)e_3) \\
&= (e_i(H^3) - H^4 \omega_3^4(e_i))(-\sin \theta e_1 - \cos \theta e_4) \\
&\quad + (e_i(H^4) + H^3 \omega_4^3(e_i))(-\sin \theta e_2 + \cos \theta e_3).
\end{aligned}$$

Comparing the above two equations, we find

$$D_{e_i}(\mathbf{J}H)^\top = (\mathbf{J}D_{e_i}^\perp H)^\top - \cos \theta (H^3 e_2 - H^4 e_1)(h_{1i}^3 + h_{2i}^4). \quad (5.6)$$

Consequently, from (5.1), (5.5) and (5.6) it follows that

$$\begin{aligned}
\operatorname{div}(\mathbf{J}H)^\top &= \sum_i \langle D_{e_i}(\mathbf{J}H)^\top, e_i \rangle \\
&= \sum_i \langle \mathbf{J}(D_{e_i}^\perp H) - \cos \theta (H^3 e_2 - H^4 e_1)(h_{1i}^3 + h_{2i}^4), e_i \rangle \\
&= \sum_i \langle \mathbf{J}(D_{e_i}^\perp H), e_i \rangle - \cos \theta (\langle (H^3 e_2 - H^4 e_1)(h_{11}^3 + h_{21}^4), e_1 \rangle \\
&\quad + \langle (H^3 e_2 - H^4 e_1)(h_{12}^3 + h_{22}^4), e_2 \rangle) \\
&= \sum_i \langle \mathbf{J}(D_{e_i}^\perp H), e_i \rangle - \cos \theta (-H^4(h_{11}^3 + h_{22}^3) + H^3(h_{11}^4 + h_{22}^4)) \\
&= \sum_i \langle \mathbf{J}(D_{e_i}^\perp H), e_i \rangle - \cos \theta (-H^4 H^3 + H^3 H^4) \\
&= - \sum_i \langle (D_{e_i}^\perp H), \mathbf{J}e_i \rangle = - \sum_i \langle h(e_i, x^\top), \mathbf{J}e_i \rangle \\
&= - \langle h(e_1, x^\top), \sin \theta e_3 \rangle - \langle h(e_2, x^\top), \sin \theta e_4 \rangle \\
&= - \langle h(e_1, x^1 e_1 + x^2 e_2), \sin \theta e_3 \rangle - \langle h(e_2, x^1 e_1 + x^2 e_2), \sin \theta e_4 \rangle \\
&= - \sin \theta (x^1 h_{11}^3 + x^2 h_{12}^3 + x^1 h_{21}^4 + x^2 h_{22}^4) \\
&= - \sin \theta (x^1 (h_{11}^3 + h_{22}^3) + x^2 (h_{11}^4 + h_{22}^4)) \\
&= - \sin \theta (x^1 H^3 + x^2 H^4) \\
&= \langle (\mathbf{J}H)^\top, x^\top \rangle,
\end{aligned}$$

where $x^\top = \sum_i x^i e_i$. If $[\alpha] = 0$, there exists a globally defined smooth function φ satisfying $\alpha = -d\varphi$, or equivalently, $\nabla \varphi = \nabla \beta_x$. By Theorem 1.2, we obtain an elliptic linear equation

$$\Delta \varphi = \frac{1}{2} \langle \nabla \varphi, \nabla |x|^2 \rangle.$$

Then according to the maximum principle (see for example, [12]), φ must be a constant and thus $\alpha = -d\varphi \equiv 0$ which with Theorem 1.2 implies $H \equiv 0$, contradicting the fact that there are no compact minimal submanifolds in an Euclidean space. $\square$

To prove Theorem 1.8 and Theorem 1.9, the following lemma is needed:

Lemma 5.1. *Let $x : M^2 \rightarrow \mathbb{C}^2$ be a self-shrinker with Kähler angle θ . Then*

$$\mathcal{L} \cos \theta = -\frac{1}{4} \cos \theta |\bar{D}\mathbf{J}_M|^2, \quad (5.7)$$

$$\frac{1}{2} \mathcal{L} \cos^2 \theta = \sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D}\mathbf{J}_M|^2. \quad (5.8)$$

Proof. Around each point of the open subset

$$M^* := \{p \in M^2; \sin \theta(p) \neq 0\},$$

we choose an orthonormal frame field $\{e_1, e_2\}$ on M^2 which via (2.2) defines the normal frame $\{e_3, e_4\}$. Then by (2.3) and (2.4)

$$\nabla \theta = \sum_i (h_{1i}^4 - h_{2i}^3) e_i, \quad \Gamma_{1i}^2 - \Gamma_{3i}^4 = \cot \theta (h_{1i}^3 + h_{2i}^4), \quad (5.9)$$

where Γ_{AB}^C are determined by $\omega_A^C = \sum_B \Gamma_{AB}^C \omega^B$. From Codazzi equation and (5.9), it follows that

$$\begin{aligned} \Delta \theta &= \sum_i (h_{1i}^4 - h_{2i}^3)_{,i} \\ &= \sum_i (e_i(h_{1i}^4 - h_{2i}^3) - (h_{1j}^4 - h_{2j}^3) \Gamma_{ii}^j) \\ &= \sum_i (h_{1ii}^4 - h_{2ii}^3 + h_{2i}^4 \Gamma_{1i}^2 - h_{1i}^3 \Gamma_{2i}^1 - h_{1i}^3 \Gamma_{3i}^4 + h_{2i}^4 \Gamma_{4i}^3) \\ &= \sum_i (H_{,1}^4 - H_{,2}^3 + (h_{2i}^4 + h_{1i}^3) \Gamma_{1i}^2 - (h_{2i}^4 + h_{1i}^3) \Gamma_{3i}^4) \\ &= \sum_i (H_{,1}^4 - H_{,2}^3 + (h_{2i}^4 + h_{1i}^3) (\Gamma_{1i}^2 - \Gamma_{3i}^4)) \\ &= \sum_i (H_{,1}^4 - H_{,2}^3 + \cot \theta (h_{2i}^4 + h_{1i}^3)^2). \end{aligned} \quad (5.10)$$

Consequently, by (5.9) and (5.10),

$$\begin{aligned} \Delta \cos \theta &= -\cos \theta |\nabla \theta|^2 - \sin \theta \Delta \theta \\ &= -\cos \theta \sum_i (h_{1i}^4 - h_{2i}^3)^2 - \sin \theta (H_{,1}^4 - H_{,2}^3) - \cos \theta \sum_i (h_{2i}^4 + h_{1i}^3)^2 \\ &= -\cos \theta \left(\sum_i ((h_{1i}^4 - h_{2i}^3)^2 + (h_{2i}^4 + h_{1i}^3)^2) \right) - \sin \theta (H_{,1}^4 - H_{,2}^3) \\ &= -\frac{1}{4} \cos \theta |\bar{D}\mathbf{J}_M|^2 - \sin \theta (H_{,1}^4 - H_{,2}^3), \end{aligned} \quad (5.11)$$

and thus

$$\begin{aligned} \frac{1}{2} \Delta (\cos^2 \theta) &= |\nabla \cos \theta|^2 + \cos \theta (\Delta \cos \theta) \\ &= \sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D}\mathbf{J}_M|^2 - \cos \theta \sin \theta (H_{,1}^4 - H_{,2}^3). \end{aligned} \quad (5.12)$$

By Lemma 2.2, we have

$$H_{,1}^4 - H_{,2}^3 = \sum_i h_{1i}^4 \langle x, e_i \rangle - \sum_i h_{2i}^3 \langle x, e_i \rangle = \sum_i (h_{1i}^4 - h_{2i}^3) \langle x, e_i \rangle. \quad (5.13)$$

Inserting (5.13) to (5.11) and (5.12), respectively, we obtain

$$\begin{aligned} \Delta \cos \theta &= -\frac{1}{4} \cos \theta |\bar{D} \mathbf{J}_M|^2 - \sin \theta \sum_i (h_{1i}^4 - h_{2i}^3) \langle x, e_i \rangle \\ &= -\frac{1}{4} \cos \theta |\bar{D} \mathbf{J}_M|^2 - \langle x, \sin \theta \sum_i (h_{1i}^4 - h_{2i}^3) e_i \rangle \\ &= -\frac{1}{4} \cos \theta |\bar{D} \mathbf{J}_M|^2 + \langle x, \nabla \cos \theta \rangle, \end{aligned} \quad (5.14)$$

and

$$\begin{aligned} \frac{1}{2} \Delta \cos^2 \theta &= |\nabla \cos \theta|^2 + \cos \theta (\Delta \cos \theta) \\ &= \sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D} \mathbf{J}_M|^2 + \cos \theta \langle x, \nabla \cos \theta \rangle \\ &= \sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D} \mathbf{J}_M|^2 + \cos \theta \langle x, \nabla \cos \theta \rangle \\ &= \sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D} \mathbf{J}_M|^2 + \frac{1}{2} \langle x, \nabla \cos^2 \theta \rangle. \end{aligned} \quad (5.15)$$

By the definition of $\mathcal{L}$, (5.14) and (5.15) become, respectively,

$$\mathcal{L} \cos \theta = \Delta \cos \theta - \langle x, \nabla \cos \theta \rangle = -\frac{1}{4} \cos \theta |\bar{D} \mathbf{J}_M|^2, \quad (5.16)$$

$$\frac{1}{2} \mathcal{L} \cos^2 \theta = \frac{1}{2} \Delta \cos^2 \theta - \frac{1}{2} \langle x, \nabla \cos^2 \theta \rangle = \sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D} \mathbf{J}_M|^2. \quad (5.17)$$

Thus (5.7) and (5.8) hold on the open set M^* .

At a point $p \notin M^*$, there are two cases that need to be considered:

Case 1°: p is an accumulation point of M^* .

Because both $\mathcal{L} \cos \theta$ and $-\cos \theta |\bar{D} \mathbf{J}_M|^2$ are globally defined and smooth, we can take the limits of the two sides of (5.7) to find (5.7) still holds at the point p . Similarly (5.8) holds at p .

Case 2°: p is not an accumulation point of M^* .

In this case, $\mathbf{J}_M = \mathbf{J}$ around p . So $|\bar{D} \mathbf{J}_M|^2 = 0$. Since $\cos \theta \equiv 1$ around p , $\mathcal{L} \cos \theta = 0$. Therefore both (5.7) and (5.8) still hold at p . $\square$

(2) *Proof of Theorem 1.8*

Take $v = \cos \theta$. Then

$$\begin{aligned} |\nabla v|^2 &= \sin^2 \theta |\nabla \theta|^2 = \sin^2 \theta \sum_i |h_{1i}^4 - h_{2i}^3|^2 \\ &\leq 2 \sin^2 \theta \sum_i ((h_{1i}^4)^2 + (h_{2i}^3)^2) \leq 2|h|^2, \end{aligned} \quad (5.18)$$

$$|\mathcal{L}v| = |\mathcal{L} \cos \theta| = \frac{1}{4} |\cos \theta| |\bar{D} \mathbf{J}_M|^2 \leq 4|h|^2. \quad (5.19)$$

Since $\int_M |h|^2 e^{-\frac{|x|^2}{2}} dV_M < \infty$, (5.18) and (5.19) show that, for $u \equiv 1$,

$$\int_M (|u \nabla v| + |\nabla u \nabla v| + |u \mathcal{L}v|) e^{-\frac{|x|^2}{2}} dV_M < \infty.$$

Thus by Lemma 2.3, we obtain

$$\int_M \mathcal{L} \cos \theta e^{-\frac{|x|^2}{2}} dV_M = 0. \quad (5.20)$$

Since, by the assumption, $\cos \theta \geq 0$, it follows by (5.7) that $\mathcal{L} \cos \theta \leq 0$. Therefore,

$$\mathcal{L} \cos \theta = -\frac{1}{4} \cos \theta |\bar{D} \mathbf{J}_M|^2 \equiv 0. \quad (5.21)$$

Consider the function $\varphi = |\nabla \theta|^2$. If φ is not identically zero on M^2 , then there exists a point p_0 such that $\varphi \neq 0$ and we can find a connected domain U containing p_0 where φ is non-zero identically. Denote $U_0 = \{p \in U \mid \cos \theta(p) = 0\}$. Then U_0 obviously contains no interior points. Therefore $U \setminus U_0$ is a dense set. By (5.21), $|\bar{D} \mathbf{J}_M|^2 \equiv 0$ on $U \setminus U_0$ implying $h_{1i}^4 = h_{2i}^3$ and $h_{2i}^4 = -h_{1i}^3$ on $U \setminus U_0$. Thus, by (2.5), we have $\nabla \theta \equiv 0$ on $U \setminus U_0$ which contradicts the definition of U . This shows that $\varphi \equiv 0$ on M^2 , that is, θ is constant.

If $\cos \theta = 0$, then x is a Lagrangian immersion;

If $\cos \theta \neq 0$, then $|\bar{D} \mathbf{J}_M|^2 \equiv 0$. It follows that $h_{1i}^4 = h_{2i}^3$, $h_{2i}^4 = -h_{1i}^3$, that is,

$$h_{11}^4 = h_{21}^3 = -h_{22}^4, \quad h_{22}^3 = h_{21}^4 = -h_{11}^3$$

which implies that $H \equiv 0$. Since x is a self-shrinker, we obtain

$$\langle x, e_3 \rangle = 0, \quad \langle x, e_4 \rangle = 0. \quad (5.22)$$

Hence, for $a = 3, 4$,

$$\langle x, \bar{D}_{e_i} e_a \rangle = e_i \langle x, e_a \rangle - \langle \bar{D}_{e_i} x, e_a \rangle = 0 - \langle e_i, e_a \rangle = 0,$$

namely

$$\langle x, -A_{e_a} e_i + \sum_b \Gamma_{ai}^b e_b \rangle = 0.$$

So that

$$\sum_j h_{ij}^a \langle x, e_j \rangle = 0, \quad a = 3, 4. \quad (5.23)$$

From (5.23), it is easily seen that

$$\begin{vmatrix} h_{11}^3 & h_{21}^3 \\ h_{12}^3 & h_{22}^3 \end{vmatrix} = \begin{vmatrix} h_{11}^4 & h_{21}^4 \\ h_{12}^4 & h_{22}^4 \end{vmatrix} = 0.$$

It follows that

$$h_{11}^3 h_{22}^3 - (h_{21}^3)^2 = h_{11}^4 h_{22}^4 - (h_{21}^4)^2 = 0$$

which with $H = 0$ shows that

$$h_{11}^3 = h_{21}^3 = h_{11}^4 = h_{21}^4 = 0.$$

That is, M^2 is totally geodesic and thus, by the completeness, M^2 must be a plane. $\square$

(3) *Proof of Theorem 1.9*

Similar to those in (2), we take $u \equiv 1$ and $v = \frac{1}{2} \cos^2 \theta$. Since

$$\begin{aligned} |\nabla v|^2 &= \cos^2 \theta \sin^2 \theta |\nabla \theta|^2 = \cos^2 \theta \sin^2 \theta \sum_i |h_{1i}^4 - h_{2i}^3|^2 \\ &\leq 2 \sin^2 \theta \sum_i ((h_{1i}^4)^2 + (h_{2i}^3)^2) \leq 2|h|^2, \end{aligned} \quad (5.24)$$

$$|\mathcal{L}v| = \frac{1}{2} |\mathcal{L} \cos^2 \theta| = |\sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D}\mathbf{J}_M|^2| \leq 6|h|^2, \quad (5.25)$$

we can use Lemma 2.3 to get

$$\frac{1}{2} \int_M (\mathcal{L} \cos^2 \theta) e^{-\frac{|x|^2}{2}} dV_M = 0. \quad (5.26)$$

On the other hand, by (1.5) and (5.8), we find

$$\begin{aligned} \frac{1}{2} \mathcal{L} \cos^2 \theta &= \sin^2 \theta |\nabla \theta|^2 - \frac{1}{4} \cos^2 \theta |\bar{D}\mathbf{J}_M|^2 \\ &\leq \sin^2 \theta \frac{\lambda \cos^2 \theta |\bar{D}\mathbf{J}_M|^2}{4(1 - \lambda \cos^2 \theta)} - \frac{1}{4} \cos^2 \theta |\bar{D}\mathbf{J}_M|^2 \\ &= - \frac{(1 - \lambda) \cos^4 \theta |\bar{D}\mathbf{J}_M|^2}{4(1 - \lambda \cos^2 \theta)} \leq 0. \end{aligned}$$

Therefore

$$\cos^4 \theta |\bar{D}\mathbf{J}_M|^2 = 0.$$

By using an argument similar to that in (2), we can conclude that either $\cos \theta \equiv 0$ or $|\bar{D}\mathbf{J}_M|^2 \equiv 0$. Then the rest of the proof is omitted here since it is the same as that of the proof of Theorem 1.8. $\square$

REFERENCES

- [1] C. Arezzo and J. Sun, Self-shrinker for the mean curvature flow in arbitrary codimension, *Math. Z.*, 274, 993-1027(2013), DOI:10.1007/S00209-012-1104-y.
- [2] A. Arsie, Maslov class and minimality in Calabi-Yau manifolds, *J. Geometry and Physics*, 35, 145-156(2000).
- [3] J. Boton, G. R. Jensen, M. Rigoli and L. M. Woodward, On conformal minimal immersions of $\mathbb{S}^2$ into $\mathbb{C}P^n$, *Math. Ann.*, 279, 599-620(1988).
- [4] H. D. Cao and H. Z. Li, A gap theorem for self-shrinkers of the mean curvature flow in arbitrary codimension, *Calc. Var. Partial Differential Equations*, 46, 879-889(2013).
- [5] I. Castro and A. M. Lerma, The clifford torus as a self-shrinker for the Lagrangian mean curvature flow, *Int. Math. Res. Not.*, 6, 1515-1527(2014).

- [6] J. Y. Chen and J. Y. Li, Mean curvature flow of surface in 4-manifolds, *Adv. Math.*, 163, 287-309(2001).
- [7] J. Y. Chen and G. Tian, Moving symplectic curves in Kähler -Einstein surfaces, *Acta Math. Sin. Engl. Ser.*, 16, 541-548(2000).
- [8] J. Y. Chen and G. Tian, Minimal surfaces in Riemannian 4-manifolds, *Geom. Funct. Anal.*, 7, 873-916(1997).
- [9] S. S. Chern and J. Wolfson, Minimal surfaces by moving frames, *A. J. Math.*, 105, 59-83(1983).
- [10] K. Cieliebak and E. Goldstein: A note on mean curvature, Maslov class and symplectic area of Lagrangian immersions, *J. Symplectic Geom.*, 2, No. 2, 261-266(2004).
- [11] T. H. Colding and W. P. Minicozzi II, Generic mean curvature flow I: generic singularities, *Ann. Math.*, 175, 755-833(2012).
- [12] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 2001.
- [13] X. L. Han and J. Sun, Translating solitons to symplectic mean curvature flows, *Ann. Glob. Anal. Geom.*, 38, 161-169(2010).
- [14] H. Z. Li and X. F. Wang, A new characterization of the clifford torus as a Lagrangian self-shrinker, *arXiv [math.DG]: 1505.05593v1*, 21, May, 2015.
- [15] H. Z. Li and Y. Wei, Classification and rigidity of self-shrinkers in the mean curvature flow, *J. Math. Soc. Japan*, 66, 709-734(2014).
- [16] X. X. Li, A classification theorem for complete minimal surfaces in S^6 with constant Kähler angles, *Arch Math.*, 72, 385-400(1999).
- [17] X. X. Li and A.-M. Li, Minimal surfaces in S^6 with constant Kähler angles, *Acta Math. Sin.*, 39, 196-203(1996).
- [18] J.-M. Morvan, Classe de Maslov d'une immersion lagrangienne et minimalité, *C. R. Acad. Sci. Paris, Série I*, 292, 633-636(1981).
- [19] Z. Q. Li, Counterexamples to the conjecture on minimal S^2 in CP^n with constant Kähler angle. *Manus. Math.*, 88, 417-431(1995).
- [20] R. Schoen and J. Wolfson, Minimizing area among Lagrangian surfaces: the mapping problem, *J. Differential Geom.*, 58, 1-86(2001).
- [21] L. M. Wood, Minimal 2-spheres in CP^n with constant Kähler angle, *Manus. Math.*, 103, 1-8(2000).

SCHOOL OF MATHEMATICS AND INFORMATION SCIENCES

HENAN NORMAL UNIVERSITY

XINXIANG 453007, HENAN

P.R. CHINA

E-mail address: xxl@henannu.edu.cn

SCHOOL OF MATHEMATICS AND INFORMATION SCIENCES

HENAN NORMAL UNIVERSITY

XINXIANG 453007, HENAN

P.R. CHINA

E-mail address: lxliliaolx@163.com